


Accepted Manuscript

Carcinogenic and non-carcinogenic health risks of metal(oid)s in tap water from Ilam city, Iran

Yadolah Fakhri, Narottam Saha, Amir Sheikhmohammadi, Milad Rasouli, Moayed Avazpour, Aziz Rahimizadeh, Seyed-Mohammad Riahi, Mansour Ghaderpoori, Hassan Keramati, Bigard Moradi, Nazak Amanidaz, Zahra Baninameh, Amin Mousavi Khaneghah



PII: S0278-6915(18)30255-2

DOI: [10.1016/j.fct.2018.04.039](https://doi.org/10.1016/j.fct.2018.04.039)

Reference: FCT 9728

To appear in: *Food and Chemical Toxicology*

Received Date: 26 November 2017

Revised Date: 8 February 2018

Accepted Date: 18 April 2018

Please cite this article as: Fakhri, Y., Saha, N., Sheikhmohammadi, A., Rasouli, M., Avazpour, M., Rahimizadeh, A., Riahi, S.-M., Ghaderpoori, M., Keramati, H., Moradi, B., Amanidaz, N., Baninameh, Z., Khaneghah, A.M., Carcinogenic and non-carcinogenic health risks of metal(oid)s in tap water from Ilam city, Iran, *Food and Chemical Toxicology* (2018), doi: 10.1016/j.fct.2018.04.039.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Carcinogenic and non-carcinogenic health risks of metal(oid)s in tap water from Ilam city, Iran

Short title: Risk assessment of metals in drinking water of Ilam city

Yadolah Fakhri¹, Narottam Saha², Amir Sheikhmohammadi³, Milad Rasouli⁴, Moayed Avazpour^{5,*}, Aziz Rahimizadeh^{6,}, Seyed-Mohammad Riahi⁷, Mansour Ghaderpoori⁸, Hassan Keramati⁹, Bigard Moradi¹⁰, Nazak Amanidaz¹¹, Zahra Baninameh¹², Amin Mousavi Khaneghah¹³**

¹Department of Environmental Health Engineering, Student Research Committee, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

²School of Earth and Environmental Sciences, The University of Queensland, Queensland, Australia.

³Department of Environmental Health Engineering, School of Public Health, Ilam University of Medical Sciences, Ilam, Iran

⁴Department of Immunology, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

⁵Department of Environmental health, University of Khoy, Khoy, Iran

⁶Food Health Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

⁷PhD in Epidemiology, department of public health , Faculty of Health, Birjand University of Medical Sciences, Birjand, Iran

⁸Department of Environmental Health Engineering, School of Health and Nutrition, Lorestan University of Medical Sciences, Khorramabad, Iran

⁹Department of Environmental Health Engineering, School of Public Health, Semnan University of Medical Sciences, Semnan, Iran.

¹⁰Department of Health Public, Kermanshah University of Medical Sciences, Kermanshah, Iran.

¹¹Environmental Health Research Center, Golestan University of Medical Sciences, Golestan, Iran.

¹²Sina Hospital, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

¹³Department of Food Science, Faculty of Food Engineering, State University of Campinas (UNICAMP), Rua Monteiro Lobato, 80. Caixa Postal: 6121, CEP: 13083-862 Campinas, Sao Paulo, Brazil.


*, **Corresponding authors:

***Moayed Avazpour**

Department of Environmental Health Engineering, School of Public Health, Ilam University of Medical Sciences, Ilam, Iran. *Email:* mo_f_1859@yahoo.com , *Phone No:* +989149768091.

****Aziz Rahimizadeh**

Food Health Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran
Telephone: +989195131958, E-mail address: rahimizadeh10@gmail.com



Carcinogenic and non-carcinogenic health risks of metal(oid)s in tap water from Ilam city, Iran

Short title: Risk assessment of metals in drinking water of Ilam city

Abstract

One of the most important pathways for exposure to metals is drinking water ingestion. Chronic or acute exposure to metals can endanger the health of the exposed population, and hence, estimation of human health risks is crucial. In the current study for the first time, the concentrations of Mercury (Hg), Arsenic (As), Zinc (Zn), Lead (Pb) and Cobalt (Co) in 120 collected tap water samples (2015, July-November) from Ilam city, Iran were investigated using flame atomic absorption spectrophotometer. Also, the metal-induced carcinogenic and non-carcinogenic risks for consumers exposed to tap drinking water were calculated. The average (range) concentrations of Hg, Zn, As, Pb and Co were defined as 0.40 ± 0.10 $\mu\text{g/L}$ (ND-0.9 $\mu\text{g/L}$), 5014 ± 5707 $\mu\text{g/L}$ (2900.00-5668.33 $\mu\text{g/L}$), 21.008 ± 2.876 $\mu\text{g/L}$ (3.5-62 $\mu\text{g/L}$), 30.38 ± 5.56 $\mu\text{g/L}$ (6-87 $\mu\text{g/L}$), and 11.34 ± 1.61 $\mu\text{g/L}$ (0.1-50 $\mu\text{g/L}$), respectively. Average concentrations of all examined metals were significantly higher than WHO and national standard recommended limits. The ranking order of metals concentrations in the tap drinking water was $\text{Zn} > \text{Pb} > \text{As} > \text{Co} > \text{Hg}$. Except for Hg and Co, at least one age group consumers were at considerable non-carcinogenic risks induced by Zn, As and Pb (Target Hazard Quotient (THQ) > 1). The rank order of age groups consumers based on THQ and Incremental lifetime cancer risk (ILCR) was $< 1 \text{ years} > 1 - 9 \text{ years} > 20 + \text{ years} > 10 - 19 \text{ years}$. The calculated Incremental lifetime cancer risk (ILCR) and cumulative carcinogenic risk (CCR) for As and Pb in all age groups were higher than 10^{-3} value. All age groups of consumers in Ilam city, especially infants ($< 1 \text{ years}$) and children (1-10 years), are at considerable non-carcinogenic and carcinogenesis risk.

Keywords: Trace metal(oid)s; tap water; risk assessment; Iran; calculated Incremental lifetime cancer risk (ILCR); Target Hazard Quotient (THQ)

1. Introduction

Economic development, unprecedented industrial revolution, and rapid population growth have raised serious concerns regarding contamination of aquatic environments by various types of contaminants. In this context, some investigations were conducted regarding the measuring of metals (Fakhri et al., 2018a; Fakhri et al., 2018b; Fakhri et al., 2017; Ghasemidehkordi et al., 2018; Shahsavani et al., 2017; Zafarzadeh et al., 2017; Farokhneshat et al., 2016; Mirzabeygi et al., 2017), polycyclic aromatic hydrocarbons (PAHs) (Hosseini et al., 2008; Karyab et al., 2016; Zhang et al., 2014), residues of drugs (Avar et al., 2016; El Fellah et al., 2017; Franquet-Griell et al., 2016; Peng et al., 2016), pesticides residue (Amirahmadi et al., 2017; Shoeibi et al., 2013; Yadollahi et al., 2012). In recent years, metals contamination of water resources has attracted global attention due to increase in their concentrations, toxic effects, non-biodegradability, highly stable and persistent characteristics among the process (Armitage et al., 2007; Sin et al., 2001; Yuan et al., 2011). Although some metals in trace amount are essential for proper functioning of living organisms, exposure to a higher concentration of them is harmful to both human and aquatic life (Shahsavani et al., 2017; Prasad, 2013; Adel et al., 2016; Dórea, 2008; Hadiani et al., 2015). Based on recommended classifications by International Agency for Research on Cancer (IARC); Zn, Cu, Mn, Cr, and Cd were regarded as non-carcinogenic metals, whereas As (group 1), Cr (group 2B), Ni (group 2B), Cd (group 1), and Co (group 2B), Pb (group 2B) were treated as potential carcinogen metals (IARC, 2004). Exposure to toxic metals results in several adverse health consequences including cardiovascular and skeletal diseases, neurotoxicity, infertility, numerous liver and kidney problems (Fakhri et al., 2017). For instance, the elevated exposure to Pb causes gastrointestinal colitis and blood

cerebral diseases (Flora et al., 2012), Cd toxicity is responsible for cardiovascular diseases and bone pain (e.g., Itai-Itai disease) (Fagerberg et al., 2017; Kobayashi, 1978), Hg poisoning impairs senses of sight, hearing and touch (Duruibe et al., 2007; Lohren et al., 2015), Cu toxicity damages kidneys and liver, and As poisoning can result in lung, skin and other cancers (IARC, 2004).

Anthropogenic activities, such as municipal, untreated industrial, and agricultural wastewater discharge, can dramatically increase the metal concentrations (Rahman et al., 2012; Saha et al., 2016; Srebotnjak et al., 2012; Su et al., 2013). Moreover, the release of toxic metals to drinking water from corroded distribution pipes can trigger metal pollution (Faier et al., 2009; Kavcar et al., 2009; Khan et al., 2015; Shanbehzadeh et al., 2014).

In recent years, estimation of carcinogenic and non-carcinogenic health risks (e.g., heart and kidney diseases) has become important, because many clinical symptoms have been observed at concentrations lower than the prescribed limits (Adel et al., 2016; Çelebi et al., 2014; Saha and Zaman, 2013), while in some cases, exceedance of prescribed levels have not caused human health problems. However, several studies have reported metal content in drinking waters in Iran (Atapour, 2012; Malakootian et al., 2014; Pirsahab et al., 2013a), but those studies are only limited to a comparison of metal concentrations with respective standard limits. Therefore, a detailed estimation of carcinogenic and non-carcinogenic health risks is critical to better inform the consumers and decision-makers regarding the metal toxicity of drinking water in Iran.

Therefore, objectives of this study were to (1) determine concentrations of five metals (Hg, As, Zn, Pb, and Co) in tap water of Ilam city, (2) compare their levels

with guideline values prescribed by various agencies and (3) estimate and compare carcinogenic and non-carcinogenic risk for different age groups of Ilam city, including < 1, 1-9, 10-19, and > 20 years old consumers.

2. Materials and Methods

2.1. Study area

Ilam city (33.6350° N, 46.4153° E), with an area of ~ 36 km², is located in the west of Iran (Figure 1). It has a population of ~194,000. Weather of this city is temperate mountainous with average precipitation of 619.5 mm/y, and temperature ranges from -13.6 to 41.2 °C (FRW, 2010).

2.2. Water sampling and pre-treatment

In a cross-sectional study in 2015 (July to November) the required sampling procedure according to grab method (Novic et al., 2017) was conducted over the course of five months. A total of 120 (6 × 20) tap water samples were collected from water distribution networks of six water resources, including Gham Gerdalan dam (GGD), Pich-e Ashoori well (PAW), Ghoch Ali wells (GAW), Gol Gol spring (GGS), Haft Cheshmeh well (HCW), and Naghlieh well (NW).

Samples were collected in pre-washed (with 20% HNO₃) plastic bottles and acidified with HNO₃ to reach a pH < 2 to prevent precipitation and adsorption of metals to the inner surface of sample holders (Buschmann et al., 2008). The collected samples were transported to Laboratory of Pharmacology (Ilam University of medical sciences) followed by their preservation at 4°C until metal analyses (Federation and Association, 2005).

2.3. Sample preparation and Apparatus

Each 10 mL water sample was digested with 5 mL HCl and 5 mL HNO₃ (Adel et al., 2016) on a hot plate until yellow color fumes ceased to evolve. The digested solution was subsequently filtered through Millipore filter paper (Whatman filter Merck, 0.45) and diluted with deionized water to a volume of 100 mL. Concentrations of As, Zn, Hg, Co, and Pb were analyzed using flame atomic absorption spectrophotometer (AAS) (BRAIC WFX-130). Moreover, hydride vapor generator (HVG) was used with flame AAS to measure As concentrations. In order to check the measurement precision, standard reference solutions with known concentrations of the analyzed metals were used. The accuracy of the method was checked by running control samples after every three samples. Each sample was measured at least three times to check the reproducibility of the measurement. Samples were reanalyzed if the relative standard deviation of the measurement exceeded 10%.

The limit of detection (LOD) for Hg, As, Zn, Pb, and Co were determined as 0.2, 0.05, 1.5, 15 and 0.15 µg/L, respectively. The Limit of quantification (LOQ) for as 0.66, 0.17, 5.25, 42, 0.49 µg/L were for Hg, As, Zn, Pb, and Co, respectively. Also, the range of precision of analysis procedure was 94% to 103%.

2.4. Estimation non-carcinogenic risk

2.4.1. Target Hazard Quotient (THQ)

The non-carcinogenic risk of metals in foods and drinking water can be estimated by quantifying THQ using Equation (1) below (Fakhri et al., 2017). THQ value >1 indicates potential non-carcinogenic health risk of the exposed population.

$$THQ = \frac{E_F \times E_D \times W_{IR} \times C}{RfD \times AT_n} \times 10^{-3} \quad (\text{Equation 1})$$

Where E_F is the exposure frequency (365 days/year), E_D is the exposure duration (average lifetime Iranian population is 70 years) (Kazemian and Tajbakhsh, 2016), W_{IR} is the water ingestion rate, which is the water consumption per kg body weight per day (mL/kg-d) according to EPA assumptions (U.S.EPA, 2004), W_{IR} for < 1, 1–10, 11–19, > 20 years age groups were considered 68, 25, 16, and 20 mL/kg-d, respectively (U.S.EPA, 2004). C is the concentration of the analyzed metals ($\mu\text{g/L}$), RfD is the oral reference dose (mg/kg-d), and $AT_n (= E_F \times E_D)$ is the average time for non-carcinogen. RfD is the highest acceptable dose that does not cause any health effect, exceedance of this limit may cause adverse health consequences (EPA, 1993). RfD for Hg, Pb, As, Zn, and Co was 0.0003, 0.0005, 0.0003, 0.3 and 0.02 mg/kg-d, respectively (EPA, 1993).

2.4.2. Total Target hazard quotient (TTHQ)

Exposure to more than one metal contaminant may cause additive and/or interactive effects, and hence, cumulative health effect from multiple metals' exposure was calculated by summing THQ value of individual metal and expressed as TTHQ as follows (Equation 2) (Mossman, 2002):

$$TTHQ = THQ_{Pb} + THQ_{As} + THQ_{Zn} + THQ_{Co} + THQ_{Hg} \quad (\text{Equation 2})$$

TTHQ value >1 indicates the probability of adverse health effects and suggests the need for undertaking a farther investigation and possible remedial action. However, TTHQ <1 represents no possible health consequence from exposure of examined metals at current consumption rate.

2.5. Estimation of carcinogenic risk

2.5.1. Estimated daily intake (EDI)

The EDI values for As and Pb were calculated using Equation (3) (Sepanlou et al., 2017):

$$EDI = W_{IR} \times C \quad (\text{Equation } 3)$$

W_{IR} is the water ingestion rate (mL / kg-d); C is the average concentrations of metals ($\mu\text{g/L}$).

2.5.2. Incremental lifetime cancer risk (ILCR)

Based on EPA human health risk models (NFPCSP, 2011), the carcinogenic risk for As and Pb was calculated using ILCR in Equation (4) (Cao et al., 2014; Sultana et al., 2017):

$$ILCR = EDI \times CSF \quad (\text{Equation } 4)$$

EDI is the estimated daily intake (mg/kg BW/day), and CSF is the cancer slope factor (Bamuwanye et al., 2015). The CSF values for Pb and As are 0.0085 and 1.5 (mg/kg-d)⁻¹, respectively (OEHHA, 2009a, b). It is worth mentioning that, CSF values are only available for Pb and As and thus, this study estimated ILCR for Pb and As. The range of acceptable cancer risk to regulatory goals is 10^{-4} to 10^{-6} (Li et al., 2013).

The cumulative carcinogenic risk (CCR) from both As and Pb was calculated by summing ILCR for As and Pb using equation (5) (Liu et al., 2013).

$$CCR = ILCR_{As} + ILCR_{Pb} \quad (\text{Equation } 5)$$

2.6. Statistical analysis

One sample T-test was performed to statistically compare the metal concentrations with safe limits prescribed. The statistical significance was considered at $p < 0.05$. All statistical tests were performed using IBM SPSS Statistics 23.0 (IBM Corporation, Armonk, NY).

3. Result and discussion

3.1. Concentration of metals

The overall ranking order of the concentration of analyzed metals in the tap water of Ilam city was $Zn > Pb > As > Co > Hg$. Additionally, the constant exposure to Hg can pose some adverse effects on the nervous, immune, and digestive systems along with kidneys, lungs, eyes, and skin (WHO, 2017). Average concentration of Hg in the tap drinking water was $0.40 \pm 0.10 \mu\text{g/L}$ (ranged from ND to $0.9 \mu\text{g/L}$) (Table 1), which is ~ 4 times higher than WHO recommended level ($0.1 \mu\text{g/L}$) (WHO, 2004) but lower than national standard limit of $0.6 \mu\text{g/L}$ (IRISI, 1996) (Figure 2). Concentrations of Hg in water samples from four water resources were below instrumental detection limit, while its concentration in samples from other two resources (GGD and GGS) were higher than WHO recommended values (Table 1).

Zn is one of the essential elements for the health of the human body in higher concentrations may cause diarrhea depression, hair loss, and poor wound healing (Deshpande et al., 2013; Strachan, 2010). Concentrations of Zn varied from $2900.00\text{--}5668.33 \mu\text{g/L}$ with an average concentration of $5014 \pm 564 \mu\text{g/L}$ (Table 1),

which is higher than both WHO guideline and national standard limit (500 µg/L) for drinking water (Figure 2) (IRISI, 1996; WHO, 2004). Based on Zn concentrations water resources are ranked as NW > PAW > HCW > GGS > GAW > GGD (Table 1).

Arsenic is one of the most toxic metalloids that originated from both anthropogenic and natural sources (Muhammad et al., 2010). Its elevated concentration in drinking water can cause melanosis, skin lesion, vascular disease, hypertension, hyperkeratosis and cancer (Ali and Tarafdar, 2003; Muhammad et al., 2010; Rahman et al., 2009). The average (\pm SD) concentration of As in the analyzed tap water was 21.008 ± 2.876 µg/L (range 3.5-62 µg/L) (Table 1). The concentration of As in tap drinking water was higher than WHO and national standard limit (10 µg/L) (Figure 2) (IRISI, 1996; WHO, 2004). Based on As concentrations, the water resources are ranked as: GGD \approx GGS > PAW > HCW > GAW > NW (Table 1).

High Pb concentrations in drinking water may be attributed to agriculture, industrial activities or domestic wastewater discharges in Ilam city. Lead is one of the most toxic metals, and infants are very sensitive to Pb toxicity. It may cause memory problems, behavioral disturbances, anemia, lung cancer, nerve damages, hypertension, stomach cancer, headache, kidney damage, asthma and abdominal pain irritability (Farkhondeh et al., 2015; Järup, 2003; Patrick, 2006; Steenland and Boffetta, 2000). The average (\pm SD) concentration of Pb in the tap water was 30.38 ± 5.56 µg/L (ranged from 6 to 87 µg/L) (Table 1), which is higher than WHO guideline (10 µg/L) but lower than national standard limit (50 µg/L) (Figure 2) (IRISI, 1996; WHO, 2004). The ranking order of water resources, based on concentrations of Pb, is GGD > GGS > PAW > GAW > NW > HCW (Table 1).

Although cobalt has been mentioned as an essential element in vitamin B12 synthesis, its high concentration can have adverse health effects, including endocrine and cardiovascular deficits and neurological (e.g., visual and hearing) impairment (Leyssens et al., 2017). The average (\pm SD) concentration of Co in the tap water was 11.34 ± 1.61 $\mu\text{g/L}$ with minimum and maximum concentrations of 0.1 $\mu\text{g/L}$ and 50 $\mu\text{g/L}$, respectively (Table 1). The concentration of Co was higher than the national standard limit (5 $\mu\text{g/L}$) (Figure 1) (IRISI, 1996). The ranking order of water resources based on Co concentrations is NW > GGD > PAW > HCW > GAW > GGS (Table 1).

Kavcar et al. (2009) indicated that corrosion of distribution water systems is an important source metal in tap drinking water (Kavcar et al., 2009). Alidadi et al., (2011) reported the concentration of Pb (11.95 ± 6.68 $\mu\text{g/L}$) in tap drinking water in Mashhad city in the northeast Iran, which was lower than our study (Alidadi et al., 2014). Pirsaeheb et al. (2013) examined metal concentrations in tap drinking water of Kermanshah city, Iran and reported that average concentrations of As, Pb, Hg, Zn, and Co was 0.328 ± 0.70 , 2.22 ± 1.97 , 0.006 ± 0.021 , 45.29 ± 72.8 , and 2.25 ± 8.22 $\mu\text{g/L}$, respectively. Their values were lower than the values found in this study (Pirsaeheb et al., 2013b). The ranking order of metals in the tap water of Kermanshah city was Zn > Co > Pb > As > Hg, which is almost similar to our study. Khan et al. (2015) reported that concentration of As, Zn, and Pb in tap drinking water of Nowshera District, Pakistan ranged from 0.01-17.5, 10-500, 20-300 $\mu\text{g/L}$, respectively. The concentration of Pb in our study is lower than Khan et al. (2015). However, concentrations of As and Zn were higher (Khan et al., 2015). Orhan Gunduz et al. (2010) presented As concentrations data in tap water in Simav plain, Turkey. Their average As concentration was 99 $\mu\text{g/L}$ and maximum concentration

was 561 $\mu\text{g/L}$, which is higher than our study (Gunduz et al., 2010). Similar to our study, Mosaferi et al. (2008) reported As concentrations higher than standard limits in the tap water of 50 villages in Hashrood, Iran (Mosaferi et al., 2008). Karim et al. (2011, examined the metal concentrations in the tap drinking water of the Karachi, Pakistan. The results of that study indicated that concentrations of Pb and Co were 6.05 ± 4.14 and 2.24 ± 1.23 $\mu\text{g/L}$, respectively, which was lower than our study. Nguyen et al. (2009), reported the average concentration of As in drinking water of three regions of Ha Nam province, Vietnam and their values (211, 348 and 325 $\mu\text{g/L}$, respectively) were higher than our study (Nguyen et al., 2009).

High concentration of metals in tap drinking water of Ilam city may be correlated to natural geologic and/or human-made sources (Fallahpour 2016). High levels of metal in groundwater of Kurdistan province, the neighboring city of Ilam, was attributed to geologic sources (Mosaferi, Yunesion et al. 2003, Ebrahimi, Eslami et al. 2015, Yousefi and Jahangard 2016). Overutilization of agrochemicals and industrial discharge may have a minor influence on higher metal concentrations in drinking water sourced from groundwater. Also, corrosion of pipes in water supply network may have significant influence in metal contamination of tap water.

3.2. Non-carcinogenic risk assessment

Except for 11–19 years old consumers, THQ values of Pb for other three groups were higher than 1 (Table 2). A striking feature of this study is that THQ values of As for all age groups exceeded the non-carcinogenic threshold value of 1, indicating that population should consume water from this area with caution. Estimated THQ

values of other metals were lower than 1, except for Zn in <1-year-old group (Table 2).

Based on THQ, the age groups are ranked as: < 1 years > 1-10 years > 20+ years > 11-19 years. Since W_{IR} for <1-year consumers was lower than others age groups, their THQ value was higher. Since W_{IR} for 11-19 year age group was lower than 20 + age groups (U.S.EPA, 2004), they had higher THQ. The contribution of metals to TTHQ is ranked as (47.9 %) > Pb (40.7 %) > Zn (11.2 %) > Hg (0.9 %) > Co (0.4 %) (Table 2, Figure 3). THQ of metals in the drinking water depends on ingestion rate, toxicity, and concentration (Kapaj et al., 2006). Similar to our study, Muhammad et al. (2010) and Khan et al. (2015) found THQ higher than 1 value in Pakistan (Kapaj et al., 2006; Muhammad et al., 2010, 2011). TTHQ in all age groups was higher than 1 (Table 2, Figure 3). Since WIR in the 11-19 years age group was lower than 20 + age groups (U.S.EPA, 2004), they had lower THQ. Since the RfD value of AS is much lower than other metals (EPA, 1993), estimated THQ for As is much higher than that of other metals, although its measured concentration was comparatively lower than other metals.

3.3. Carcinogenic risk assessment

As, and Pb are categorized as carcinogens by the IARC (IARC, 2002; Fakhri et al., 2015; Farokhi et al., 2017; Khelifi et al., 2013). Since CSF value for Cd is not available, we excluded Cd from quantifying ILCR. The minimum and maximum ILCR of As and Pb was observed in the 10–19 (1.99E-01 and 1.63E-03 for As and Pb, respectively) and < 1 (8.44E-01 and 6.92E-03 for As and Pb, respectively) year age group consumers, respectively (Table 3). The cancer risk of As in the all age groups

were 121 times higher than Pb. Since W_{IR} in the < 1 years consumers were higher than the other age groups, the ILCR was high in them (EPA, 2017; USEPA, 2000, 2016). When $ILCR > 10^{-3}$, $ILCR > 10^{-4}$, and $ILCR < 10^{-6}$ consumers are in the considerable risk range, threshold risk range and safe range, respectively (EPA, 2017; USEPA, 2000, 2016). ILCR for all age group consumers from both As and Pb were $> 10^{-3}$ (Table 3). Moreover, the CCR values for As and Pb were $> 10^{-3}$ (Table 4, Figure 4). The ranking order of the age groups based on CCR was < 1 years > 1-9 years > + 20 years > 11-19 years (Figure 5). Similar to our study, CR of As in Khan et al. (2015) (ranged from 1.5E-09 to 1.3E+06) and Nguyen et al. (2009) (average 4E-4) was higher than 10^{-6} (Khan et al., 2015; Nguyen et al., 2009).

4. Conclusions

In the current study for the first time, the concentrations of Mercury (Hg), Arsenic (As), Zinc (Zn), Lead (Pb) and Cobalt (Co) in 120 collected tap water samples (2015 July-November) from Ilam city, Iran were investigated using flame atomic absorption spectrophotometer. Also, the metal-induced carcinogenic and non-carcinogenic risks for consumers exposed to tap drinking water were calculated. Average concentrations of all metals investigated (Zn, Pb, As, Co, and Hg) in tap drinking water of Ilam city were significantly higher than WHO and national standard recommended limits. Moreover, the rank order of metals based on quota in the TTHQ was As > Pb > Zn > Hg > Co. All age consumers are at considerable non-carcinogenic risk of metals in the tap drinking water ($TTHQ > 1$). Additionally, all age consumers are at considerable carcinogenic risk of As and Pb in the tap drinking water of Ilam ($ILCR > 10^{-3}$; $CCR > 10^{-3}$). CR risk of As was ~ 122 times higher than

Pb. Therefore, further considerations should be given to reduce the concentration of As in the drinking water resources. Generally speaking, the results of this study indicated that drinking water consumers in Ilam, especially infants (< 1 years) and children (1-10 years), are at serious non-carcinogenicity and carcinogenesis risks. To minimize the health threats of the local population of Ilam city, appropriate water treatment policy should be implemented to ensure safe water supply. Integrated water management approach should be employed to prevent contamination of water supply system and to protect the source water from anthropogenic metal pollution, which can be achieved by identifying the uses of chemicals through regular inspection of the catchment areas.

Acknowledgement

The authors would like to thank student research committee, Ilam University of medical sciences for the financial grants of this study (code 938045/150). Amin Mousavi Khaneghah gratefully acknowledges the support of CNPq-TWAS Postgraduate Fellowship (Grant #3240274290).

Declaration of Interest

There is no conflict of interest.

References

Adel, M., Dadar, M., Fakhri, Y., Oliveri Conti, G., Ferrante, M., 2016. Heavy metal concentration in muscle of pike (*Esox lucius* Linnaeus, 1758) from Anzali international wetland, southwest of the Caspian Sea and their consumption risk assessment. *Toxin Reviews* 35, 217-223.

- 1 Ali, M., Tarafdar, S., 2003. Arsenic in drinking water and in scalp hair by EDXRF: A
2 major recent health hazard in Bangladesh. *Journal of Radioanalytical and Nuclear*
3 *Chemistry* 256, 297-305.
- 4 Alidadi, H., Peiravi, R., Dehghan, A.A., Vahedian, M., Moalemzade Haghighi, H.,
5 Amini, A., 2014. Survey of heavy metals concentration in Mashhad drinking water in
6 2011. *Razi Journal of Medical Sciences* 20, 27-34.
- 7 Amirahmadi, M., Kobarfard, F., Pirali-Hamedani, M., Yazdanpanah, H., Rastegar, H.,
8 Shoeibi, S., Mousavi Khaneghah, A., Effect of Iranian traditional cooking on fate of
9 pesticides in white rice. *Toxin Revi* 36, 2017, 177-186.
- 10 Armitage, P.D., Bowes, M.J., Vincent, H.M., 2007. Long-term changes in
11 macroinvertebrate communities of a heavy metal polluted stream: the river Nent
12 (Cumbria, UK) after 28 years. *River research and applications* 23, 997-1015.
- 13 Atapour, H., 2012. Geochemical baseline of major anions and heavy metals in
14 groundwaters and drinking waters around the urban areas of Kerman city,
15 southeastern Iran. *Environmental Earth Sciences* 67, 2063-2076.
- 16 Avar, P., Maász, G., Takács, P., Lovas, S., Zrínyi, Z., Svigruha, R., Takátsy, A.,
17 Tóth, L., Pirger, Z., 2016. HPLC-MS/MS analysis of steroid hormones in
18 environmental water samples. *Drug testing and analysis* 8, 123-127.
- 19 Bamuwamye, M., Ogwok, P., Tumuhairwe, V., 2015. Cancer and non-cancer risks
20 associated with heavy metal exposures from street foods: Evaluation of roasted
21 meats in an urban setting. *Journal of Environment Pollution and Human Health* 3,
22 24-30.
- 23 Buschmann, J., Berg, M., Stengel, C., Winkel, L., Sampson, M.L., Trang, P.T.K.,
24 Viet, P.H., 2008. Contamination of drinking water resources in the Mekong delta

- 1 floodplains: Arsenic and other trace metals pose serious health risks to population.
2 Environment International 34, 756-764.
- 3 Cao, S., Duan, X., Zhao, X., Ma, J., Dong, T., Huang, N., Sun, C., He, B., Wei, F.,
4 2014. Health risks from the exposure of children to As, Se, Pb and other heavy
5 metals near the largest coking plant in China. Science of the Total Environment 472,
6 1001-1009.
- 7 Çelebi, A., Şengörür, B., Kløve, B., 2014. Human health risk assessment of
8 dissolved metals in groundwater and surface waters in the Melen watershed, Turkey.
9 Journal of Environmental Science and Health, Part A 49, 153-161.
- 10 Deshpande, J.D., Joshi, M.M., Giri, P.A., 2013. Zinc: The trace element of major
11 importance in human nutrition and health.
- 12 Dórea, J.G., 2008. Persistent, bioaccumulative and toxic substances in fish: human
13 health considerations. Science of the Total Environment 400, 93-114.
- 14 Duruibe, J., Ogwuegbu, M., Ekwurugwu, J., 2007. Heavy metal pollution and human
15 biotoxic effects. International Journal of Physical Sciences 2, 112-118.
- 16 Ebrahimi, S. J. A., A. Eslami and L. Ebrahimzadeh., 2015. Evaluation of Heavy
17 Metals Concentration in the Drinking Water Distribution Network in Kurdistan
18 Villages in the Year 2012. Research Journal Of Pharmaceutical Biological And
19 Chemical Sciences, 6(2), 55-61.
- 20 El Fellah, S., Duporté, G., Sirén, H., 2017. Steroid hormones, inorganic ions and
21 botrydial in drinking water. Determination with capillary electrophoresis and liquid
22 chromatography-orbitrap high resolution mass spectrometry. Microchemical Journal
23 133, 126-136.

- 1 EPA, 1993. Reference Dose (RfD): Description and Use in Health Risk
2 Assessments, [https://www.epa.gov/iris/reference-dose-rfd-description-and-use-](https://www.epa.gov/iris/reference-dose-rfd-description-and-use-health-risk-assessments)
3 [health-risk-assessments](https://www.epa.gov/iris/reference-dose-rfd-description-and-use-health-risk-assessments).
- 4 EPA, 2017. Regional Screening Levels (RSLs) - Generic Tables (June
5 2017)[https://www.epa.gov/risk/regional-screening-levels-rsls-generic-tables-june-](https://www.epa.gov/risk/regional-screening-levels-rsls-generic-tables-june-2017)
6 2017.
- 7 Fagerberg, B., Borné, Y., Barregard, L., Sallsten, G., Forsgard, N., Hedblad, B.,
8 Persson, M., Engström, G., 2017. Cadmium exposure is associated with soluble
9 urokinase plasminogen activator receptor, a circulating marker of inflammation and
10 future cardiovascular disease. *Environmental Research* 152, 185-191.
- 11 Fallahpour, N. 2016. Effect of natural organic matter, metal ions, and nitrate on
12 electrochemical dechlorination of trichloroethylene, Northeastern University.
- 13 Faier, M.C., Dumitre, G., Perju, D., 2009. Experimental Modeling of Heavy Metals
14 Concentration Distribution in Rivers.
- 15 Fakhri, Y., Mousavi Khaneghah, A., Hadiani, M.R., Keramati, H., Hosseini Pouya, R.,
16 Moradi, B., da Silva, B.S., Non-carcinogenic risk assessment induced by heavy
17 metals content of the bottled water in Iran. *Toxin Revi* 36, 2017, 313-321.
- 18 Fakhri, Y., Mohseni-Bandpei, A., Oliveri Conti, G., Keramati, H., Zandsalimi, Y.,
19 Amanidaz, N., Hosseini Pouya, R., Moradi, B., Bahmani, Z., Rasouli Amirhajloo, L.,
20 2017. Health risk assessment induced by chloroform content of the drinking water in
21 Iran: systematic review. *Toxin Reviews* 36, 1-11.
- 22 Fakhri, Y., Mohseni-Bandpei, A., Conti, G.O., Ferrante, M., Cristaldi, A., Jeihooni,
23 A.K., Alinejad, A., Mohseni, S.M., Sarkhosh, M., Keramati, H., Systematic review
24 and health risk assessment of arsenic and lead in the fished shrimps from the
25 Persian gulf. *Food and Chemical Toxicology*, 2018b.

- 1
- 2 Fakhri, Y., Mohseni, S.M., Jafarzadeh, S., Langarizadeh, G., Moradi, B., Zandsalimi,
- 3 Y., Rahimizadeh, A., Mirzaei, M., 2015. Assessment of carcinogenic and non-
- 4 carcinogenic risk lead in bottled water in different age groups in Bandar Abbas City,
- 5 Iran. *Global journal of health science* 7, 286.
- 6 Fakhri, Y., Mohseni-Bandpei, A., Oliveri Conti, G., Keramati, H., Zandsalimi, Y., Amanidaz,
- 7 N., Hosseini Pouya, R., Moradi, B., Bahmani, Z., Rasouli Amirhajloo, L., Health risk
- 8 assessment induced by chloroform content of the drinking water in Iran: systematic review.
- 9 *Toxin Revi* 36, 2017, 322-330.
- 10 Farkhondeh, T., Samarghandian, S., Sadighara, P., 2015. Lead exposure and
- 11 asthma: an overview of observational and experimental studies. *Toxin Reviews* 34,
- 12 6-10.
- 13 Farokhi, A., Tayebipoor, M., Koohpeima, F., Mokhtari, M.J., 2017. Lead and cancer.
- 14 *Journal of Cellular Immunotherapy* 3, 2.
- 15 Farokhneshat, F., Rahmani, A., Samadi, M., Soltanian, A., 2016. Non-Carcinogenic
- 16 Risk Assessment of Heavy Metal of Lead, Chro-mium and Zinc in Drinking Water
- 17 Supplies of Hamadan in Winter 2015. *Scientific Journal of Hamadan University of*
- 18 *Medical Sciences* 23, 25-33.
- 19 Federation, W.E., Association, A.P.H., 2005. Standard methods for the examination
- 20 of water and wastewater. American Public Health Association (APHA): Washington,
- 21 DC, USA.
- 22 Flora, G., Gupta, D., Tiwari, A., 2012. Toxicity of lead: a review with recent updates.
- 23 *Interdiscip Toxicol* 5, 47-58.

- 1 Franquet-Griell, H., Ventura, F., Boleda, M.R., Lacorte, S., 2016. Do cytostatic drugs
2 reach drinking water? The case of mycophenolic acid. *Environmental Pollution* 208,
3 532-536.
- 4 FRW, 2010. Wheather of
5 Ilam.<http://ilam.frw.org.ir/00/Fa/StaticPages/Page.aspx?tid=1691>.
- 6 Ghasemidehkordi, B., Malekirad, A.A., Nazem, H., Fazilati, M., Salavati, H., Shariatifar, N.,
7 Rezaei, M., Khaneghah, A.M., Fakhri, Y., Concentration of lead and mercury in collected
8 vegetables and herbs from Markazi province, Iran: Non-carcinogenic risk assessment. *Food*
9 *and Chemical Toxicology*, 2018.
- 10 Gunduz, O., Simsek, C., Hasozbek, A., 2010. Arsenic pollution in the groundwater of
11 Simav Plain, Turkey: its impact on water quality and human health. *Water, air, and*
12 *soil pollution* 205, 43.
- 13 Hosseini, S.V., Behrooz, R.D., Esmaili-Sari, A., Bahramifar, N., Hosseini, S.M.,
14 Tahergorabi, R., Hosseini, S.F., Feás, X., 2008. Contamination by organochlorine
15 compounds in the edible tissue of four sturgeon species from the Caspian Sea (Iran).
16 *Chemosphere* 73, 972-979.
- 17 International Agency for Research on Cancer (IARC), 2002. International Agency for
18 Research on Cancer, handbooks of cancer prevention. The Agency.
- 19 International Agency for Research on Cancer (IARC), 2004. IARC Monographs on
20 the
21 Evaluation of Carcinogenic Risk to Human. Some Drinking-water Disinfectants
22 and Contaminants, Including arsenic, Lyons, France, 84, 39–270.
- 23 Iran of Research Industrial and Standards of Institute (IRISI), 1996. Drinking water
24 Physical and chemical specifications , ISIRI :1053 , revision.5th, www.isiri.gov.ir/.

- 1 Järup, L., 2003. Hazards of heavy metal contamination. British medical bulletin 68,
2 167-182.
- 3 Kapaj, S., Peterson, H., Liber, K., Bhattacharya, P., 2006. Human health effects from
4 chronic arsenic poisoning—a review. Journal of Environmental Science and Health
5 Part A 41, 2399-2428.
- 6 Karim, Z., 2011. Risk assessment of dissolved trace metals in drinking water of
7 Karachi, Pakistan. Bulletin of environmental contamination and toxicology 86, 676-
8 678.
- 9 Karyab, H., Yunesian, M., Nasser, S., Rastkari, N., Mahvi, A., Nabizadeh, R., 2016.
10 Carcinogen Risk Assessment of Polycyclic Aromatic Hydrocarbons in Drinking
11 Water, Using Probabilistic Approaches. Iranian journal of public health 45, 1455.
- 12 Kavcar, P., Sofuoglu, A., Sofuoglu, S.C., 2009. A health risk assessment for
13 exposure to trace metals via drinking water ingestion pathway. International journal
14 of hygiene and environmental health 212, 216-227.
- 15 Kazemian, M., Tajbakhsh, E., 2016. Revising life expectancy index to explain its
16 consistency to other vital indices in Iran.
- 17 Khan, S., Shah, I.A., Muhammad, S., Malik, R.N., Shah, M.T., 2015. Arsenic and
18 heavy metal concentrations in drinking water in Pakistan and risk assessment: a
19 case study. Human and Ecological Risk Assessment: An International Journal 21,
20 1020-1031.
- 21 Khelifi, R., Olmedo, P., Gil, F., Hammami, B., Chakroun, A., Rebai, A., Hamza-
22 Chaffai, A., 2013. Arsenic, cadmium, chromium and nickel in cancerous and healthy
23 tissues from patients with head and neck cancer. Science of the Total Environment
24 452, 58-67.

- 1 Kobayashi, J., 1978. Pollution by cadmium and the itai-itai disease in Japan. Toxicity
2 of heavy metals in the environment, 199-260.
- 3 Koedrith, P., Seo, Y.R., 2011. Advances in carcinogenic metal toxicity and potential
4 molecular markers. International journal of molecular sciences 12, 9576-9595.
- 5 Leyssens, L., Vinck, B., Van Der Straeten, C., Wuyts, F., Maes, L., 2017. Cobalt
6 toxicity in humans—A review of the potential sources and systemic health effects.
7 Toxicology 387, 43-56.
- 8 Li, P.-H., Kong, S.-F., Geng, C.-M., Han, B., Lu, B., Sun, R.-F., Zhao, R.-J., Bai, Z.-
9 P., 2013. Assessing the hazardous risks of vehicle inspection workers' exposure to
10 particulate heavy metals in their work places. Aerosol and Air Quality Research 13,
11 255-265.
- 12 Liu, X., Song, Q., Tang, Y., Li, W., Xu, J., Wu, J., Wang, F., Brookes, P.C., 2013.
13 Human health risk assessment of heavy metals in soil–vegetable system: a multi-
14 medium analysis. Science of the Total Environment 463, 530-540.
- 15 Lohren, H., Blagojevic, L., Fitkau, R., Ebert, F., Schildknecht, S., Leist, M.,
16 Schwerdtle, T., 2015. Toxicity of organic and inorganic mercury species in
17 differentiated human neurons and human astrocytes. Journal of Trace Elements in
18 Medicine and Biology 32, 200-208.
- 19 Hadiani, M.R., Dezfooli-Manesh, S., Shoeibi, S., Ziarati, P., Mousavi Khaneghah, A.,
20 Trace elements and heavy metals in mineral and bottled drinking waters on the
21 Iranian market. Food Additives & Contaminants: Part B 8, 2015, 18-24.
- 22 Malakootian, M., Mobini, M., Sharife, I., 2014. Evaluation of Corrosion and Scaling
23 Potential of Wells Drinking Water and Aqueducts in Rural Areas Adjacent to
24 Rafsanjan Fault in During October to December 2013. Journal of Rafsanjan
25 University of Medical Sciences 13, 293-304.

- 1 Mirzabeygi, M., Abbasnia, A., Yunesian, M., Nodehi, R.N., Yousefi, N., Hadi, M.,
- 2 Mahvi, A.H., 2017. Heavy metal contamination and health risk assessment in
- 3 drinking water of Sistan and Baluchistan, Southeastern Iran. *Human and Ecological*
- 4 *Risk Assessment: An International Journal* 23, 1893-1905.
- 5 Mosaferi, M., Taghipour, H., Hassani, A., Borghei, M., Kamali, Z., Ghadirzadeh, A.,
- 6 2008. Study of arsenic presence in drinking water sources: a case study. *Iranian*
- 7 *Journal of Health and Environment* 1, 19-28.
- 8 Mosaferi, M., et al. 2003. Arsenic occurrence in drinking water of IR of Iran: the case
- 9 of Kurdistan Province. *Fate of arsenic in the environment*. Dhaka: BUET-UNU
- 10 International Symposium, International Training Network Centre, Bangladesh
- 11 University of Engineering and Technology, United Nations University, Tokyo.
- 12 Mossman, K.L., 2002. *Human and ecological risk assessment: Theory and practice*.
- 13 LWW.
- 14 Muhammad, S., Shah, M.T., Khan, S., 2010. Arsenic health risk assessment in
- 15 drinking water and source apportionment using multivariate statistical techniques in
- 16 Kohistan region, northern Pakistan. *Food and Chemical Toxicology* 48, 2855-2864.
- 17 Muhammad, S., Shah, M.T., Khan, S., 2011. Health risk assessment of heavy metals
- 18 and their source apportionment in drinking water of Kohistan region, northern
- 19 Pakistan. *Microchemical Journal* 98, 334-343.
- 20 NFPCSP, 2011. *Nutrition Fact Sheet*. Joint report of Food Planning and Nutrition
- 21 Unit (FMPU) of the ministry of Food of Government of Bangladesh and Food and
- 22 Agricultural Organization of the United Nation (FAO), September 14, 1–2. *National*
- 23 *Food Policy Plan of Action and Country Investment Plan*, Government of the
- 24 *People's Republic of Bangladesh*.

- 1 Nguyen, V.A., Bang, S., Viet, P.H., Kim, K.-W., 2009. Contamination of groundwater
2 and risk assessment for arsenic exposure in Ha Nam province, Vietnam.
3 Environment International 35, 466-472.
- 4 Novic, A.J., O'Brien, D.S., Kaserzon, S.L., Hawker, D.W., Lewis, S.E., Mueller, J.F.,
5 2017. Monitoring Herbicide Concentrations and Loads during a Flood Event: A
6 Comparison of Grab Sampling with Passive Sampling. Environmental Science &
7 Technology 51, 3880-3891.
- 8 OEHHA, 2009a. Air Toxics Hot Spots Program Technical Support Document for
9 Cancer Potencies. Appendix B. Chemical-specific summaries of the information used
10 to derive unit risk and cancer potency values. Updated 2011.
- 11 OEHHA, 2009b. Office of Environmental Health Hazard Assessment. Lead and Lead
12 Compounds. <https://oehha.ca.gov/chemicals/lead-and-lead-compounds>.
- 13 Patrick, L., 2006. Lead toxicity, a review of the literature. Part I: exposure, evaluation,
14 and treatment. Alternative medicine review 11, 2-23.
- 15 Peng, Y., Hall, S., Gautam, L., 2016. Drugs of abuse in drinking water—a review of
16 current detection methods, occurrence, elimination and health risks. TrAC Trends in
17 Analytical Chemistry 85, 232-240.
- 18 Pirsaeheb, M., Dargahi, A., Golestanifar, H., 2013a. Determination of arsenic in
19 agricultural products, animal products and drinking water of rural areas of Bijar and
20 Gharve, Kurdistan Province.
- 21 Pirsaeheb, M., Khosravi, T., Sharafi, K., Babajani, L., Rezaei, M., 2013b.
22 Measurement of heavy metals concentration in drinking water from source to
23 consumption site in Kermanshah-Iran. World Applied Sciences Journal 21, 416-423.

- 1 Rahman, M.M., Naidu, R., Bhattacharya, P., 2009. Arsenic contamination in
2 groundwater in the Southeast Asia region. *Environmental Geochemistry and Health*
3 31, 9-21.
- 4 Rahman, M.S., Molla, A.H., Saha, N., Rahman, A., 2012. Study on heavy metals
5 levels and its risk assessment in some edible fishes from Bangshi River, Savar,
6 Dhaka, Bangladesh. *Food Chemistry* 134, 1847-1854.
- 7 Shoeibi, S., Amirahmadi, M., Rastegar, H., Khosrokhavar, R., Khaneghah, A.M., An
8 applicable strategy for improvement recovery in simultaneous analysis of 20
9 pesticides residue in tea. *J Food Sci* 78, 2013.
- 10 Saha, N., Mollah, M., Alam, M., Rahman, M.S., 2016. Seasonal investigation of
11 heavy metals in marine fishes captured from the Bay of Bengal and the implications
12 for human health risk assessment. *Food control* 70, 110-118.
- 13 Saha, N., Zaman, M., 2013. Evaluation of possible health risks of heavy metals by
14 consumption of foodstuffs available in the central market of Rajshahi City,
15 Bangladesh. *Environmental monitoring and assessment* 185, 3867-3878.
- 16 Sepanlou, S.G., Parsaeian, M., Krohn, K.J., Afshin, A., Farzadfar, F., Roshandel, G.,
17 Karimkhani, C., Bazargan-Hejazi, S., Kiadaliri, A.A., Ahmadi, H., 2017. Disability-
18 adjusted life-years (DALYs) for 315 diseases and injuries and healthy life expectancy
19 (hale) in Iran and its neighboring countries, 1990-2015: findings from Global Burden
20 of Disease Study 2015. *Archives of Iranian medicine* 20, 403-418.
- 21 Shanbehzadeh, S., Vahid Dastjerdi, M., Hassanzadeh, A., Kiyanizadeh, T., 2014.
22 Heavy metals in water and sediment: a case study of Tembi River. *Journal of*
23 *environmental and public health* 2014.

- 1 Shahsavani, A., Fakhri, Y., Ferrante, M., Keramati, H., Zandsalimi, Y., Bay, A., Hosseini
2 Pouya, S.R., Moradi, B., Bahmani, Z., Mousavi Khaneghah, A., Risk assessment of heavy
3 metals bioaccumulation: fished shrimps from the Persian Gulf. *Toxin Rev* 36, 2017, 322-330
- 4 Sin, S., Chua, H., Lo, W., Ng, L., 2001. Assessment of heavy metal cations in
5 sediments of Shing Mun River, Hong Kong. *Environment international* 26, 297-301.
- 6 Srebotnjak, T., Carr, G., de Sherbinin, A., Rickwood, C., 2012. A global Water
7 Quality Index and hot-deck imputation of missing data. *Ecological Indicators* 17, 108-
8 119.
- 9 Shoeibi, S., Amirahmadi, M., Rastegar, H., Khosrokhavar, R., Khaneghah, A.M., An
10 applicable strategy for improvement recovery in simultaneous analysis of 20 pesticides
11 residue in tea. *J Food Sci* 78, 2013.
- 12 Steenland, K., Boffetta, P., 2000. Lead and cancer in humans: where are we now?
13 *American journal of industrial medicine* 38, 295-299.
- 14 Strachan, S., 2010. Heavy metal. *Curr. Anaesth. Crit. Care* 21, 44-48.
- 15 Su, S., Xiao, R., Mi, X., Xu, X., Zhang, Z., Wu, J., 2013. Spatial determinants of
16 hazardous chemicals in surface water of Qiantang River, China. *Ecological*
17 *indicators* 24, 375-381.
- 18 Sultana, M.S., Rana, S., Yamazaki, S., Aono, T., Yoshida, S., 2017. Health risk
19 assessment for carcinogenic and non-carcinogenic heavy metal exposures from
20 vegetables and fruits of Bangladesh. *Cogent Environmental Science* 3, 1291107.
- 21 U.S. Environmental Protection Agency (U.S. EPA), 2004. Estimated per capita water
22 ingestion and body weight in the United States -an update. 312-314.
- 23 U.S. Environmental Protection Agency (U.S. EPA), 2000. Guidance for assessing
24 chemical contaminant data for use in fish advisories, volume II. Risk Assessment

and fish consumption limits. (EPA 823-B-00-008). United States Environmental Protection Agency, Washington, DC.

U.S. Environmental Protection Agency (U.S. EPA), 2016.. Integrated Risk Information System. <https://www.epa.gov/iris/> (accessed 14.10.16).

World Health Organization (WHO), 2004. Guidelines for drinking-water quality: recommendations..

World Health Organization (WHO), 2017. Mercury and health. <http://www.who.int/mediacentre/factsheets/fs361/en/>.

Yadolahi, M., Babri, M., Sharif, A.A.M., Khaneghah, A.M., Pesticide residue determination in Shahr-e-Rey tomatoes using Quechers method. *Advances in Environmental Biology* 6, 2012, 2434-2438.

Yuan, G.-L., Liu, C., Chen, L., Yang, Z., 2011. Inputting history of heavy metals into the inland lake recorded in sediment profiles: Poyang Lake in China. *Journal of hazardous materials* 185, 336-345.

Zafarzadeh, A., Bay, A., Fakhri, Y., Keramati, H., Hosseini Pouya, R., Heavy metal (Pb, Cu, Zn, and Cd) concentrations in the water and muscle of common carp (*Cyprinus carpio*) fish and associated non-carcinogenic risk. *Toxin Revi* 36, 2017, 331-341.

Zhang, J., Li, Y., Wang, Y., Zhang, Y., Zhang, D., Zhang, R., Li, J., Zhang, G., 2014. Spatial distribution and ecological risk of polychlorinated biphenyls in sediments from Qinzhou Bay, Beibu Gulf of South China. *Marine pollution bulletin* 80, 338-343.

Table captions

Table 1. Concentrations of metals ($\mu\text{g/L}$) in the tap drinking water related to water resources supply of Ilam city.

Table 2. Nun-carcinogenic risk induced by metals in tap drinking water in the age groups consumers of Ilam city.

Table 3. Cancer risk induced by As and Pb in drinking water in the age groups consumers of Ilam city.

Table 4. CCR induced by As and Pb in drinking water in the age groups consumers of Ilam city.

Figure captions

Figure 1. Locations of water reserviours in the Ilam city.

Figure 2. Compare concentrations of metals in the tap drinking water with WHO guideline and national standard limits.

Figure 3. TTHQ of metals (Pb, As, Hg, Zn, and Co) in the all age groups consumers in Ilam city.

Figure 4. The contribution of each metal in the calculated TTHQ for drinking water

Figure 5. Cumulative carcinogenic risk in the age groups induced by As and Pb in the tap drinking water od Ilam city.

Table 1. Concentrations of heavy metal ($\mu\text{g/L}$) in the tap drinking water related to water resources supply of Ilam city

Heavy metals	Water resources	Sample size	Average (SD)	Minimum	Maximum
Hg	Gham Gerdalan dam	20	0.605 ± 0.19	0.30	0.90
	Pich Ashoori well	20	ND ¹	ND	ND
	Haf Cheshmeh well	20	ND	ND	ND
	Ghoch Ali wells	20	ND	ND	ND
	Naghlieh well	20	ND	ND	ND
	Gol Gol spring	20	0.21 ± 0.02	ND	0.90
	Average		0.40 ± 0.10	ND	0.90
Zn	Gham Gerdalan dam	20	4404 ± 1022	2900.00	5800.00
	Pich Ashoori well	20	5094 ± 206	4800.00	5500.00
	Haf Cheshmeh well	20	5074 ± 493	4700.00	5710.00
	Ghoch Ali wells	20	4694 ± 396	4000.00	5600.00
	Naghlieh well	20	6044 ± 680	4500.00	5800.00
	Gol Gol spring	20	4794 ± 622	3600.00	5600.00
	Average		5014 ± 564	2900.00	5668.33
As	Gham Gerdalan dam	20	50.75 ± 7.15	35.00	62.00
	Pich Ashoori well	20	9.50 ± 1.53	7.00	12.00
	Haf Cheshmeh well	20	6.20 ± 0.34	5.80	7.00
	Ghoch Ali wells	20	5.10 ± 0.74	3.50	6.10
	Naghlieh well	20	3.80 ± 0.20	30.00	4.20
	Gol Gol spring	20	50.70 ± 7.29	3.50	36.00
	Average		21.00 ± 2.87	3.50	62.00
Pb	Gham Gerdalan dam	20	67.70 ± 10.67	50.00	85.00
	Pich Ashoori well	20	16.88 ± 1.16	15.00	18.60
	Haf Cheshmeh well	20	7.50 ± 0.82	6.00	8.90
	Ghoch Ali wells	20	15.91 ± 3.53	10.00	22.00
	Naghlieh well	20	7.89 ± 0.91	6.00	8.90
	Gol Gol spring	20	66.40 ± 16.82	45.00	87.00
	Average		30.38 ± 5.56	6.00	87.00
Co	Gham Gerdalan dam	20	14.12 ± 4.07	9.00	20.00
	Pich Ashoori well	20	8.90 ± 0.83	7.00	10.00
	Haf Cheshmeh well	20	1.89 ± 0.16	1.30	2.20
	Ghoch Ali wells	20	0.45 ± 0.08	0.30	0.60
	Naghlieh well	20	42.30 ± 4.35	35.00	50.00
	Gol Gol spring	20	0.37 ± 0.16	0.10	0.70
	Average		11.34 ± 1.61	0.10	50.00

¹ Not detected

Table 2. Non-carcinogenic risk of heavy metals in population of different age groups of Ilam city.

Age consumer	THQ					TTHQ
	Pb	As	Hg	Co	Zn	
< 1	4.131	4.760	0.090	0.038	1.136	10.157
1-10	1.519	1.750	0.033	0.014	0.417	3.734
11-19	0.972	1.120	0.021	0.009	0.267	2.390
> 20	1.215	1.400	0.026	0.011	0.334	2.987
All age	1.960	2.258	0.043	0.018	0.539	4.817

Table 3. Estimated cancer risk induced by As and Pb in drinking water consumed by different age groups of Ilam city.

	Age Groups	EDI (mg/kg-d)	ILCR
As	< 1	0.00143	8.44E-01
	1-9	0.00053	3.10E-01
	10-19	0.00034	1.99E-01
	20 +	0.00042	2.48E-01
Pb	< 1	0.00207	6.92E-03
	1-9	0.00076	2.54E-03
	10-19	0.00049	1.63E-03
	20 +	0.00061	2.04E-03

Table 4. CCR induced by As and Pb in drinking water in the age group consumers of Ilam city.

Age Groups (Years)	< 1	1-10	11-19	20 +
CCR	8.51E-01	3.13E-01	2.00E-01	2.50E-01



Figure 1. Locations of water reservoirs in Ilam city, Iran.




Figure 2. Comparison of metal concentrations in tap drinking water of Ilam city with WHO guideline and national standard limits.




Figure 3 . TTHQ of metals (Pb, As, Hg, Zn, and Co) for consumers of different age groups in Ilam city.




Figure 4. The contribution of each metal in the calculated TTHQ for drinking water




Figure 5. Cumulative carcinogenic risk in four age groups induced by As and Pb in the tap drinking water of Ilam city.

- Mean concentration of metals was higher than WHO and national standard limits.
- Populations are exposed to carcinogenic and non-carcinogenic health risks.
- Infants and children are at higher risk relative to adults.